

Immune System Response after Immobilization Stress in the Background of Ionizing Radiation

Oralbek Ilderbayev^{1,*}, Gulmira Zharmakhanova²,
Saule Rakhyzhanova³, Assem Musaynova³, Gulzhan Ilderbayeva⁴,
Akmaral Nursafina¹ and Lashyn Zhazykbayeva³

¹Department of General Biology and Genomics, Faculty of Natural Sciences, L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan

²Department of Natural Science Disciplines, West Kazakhstan Marat Ospanov Medical University, Aktobe, Kazakhstan

³Department of Microbiology, Physiology, Immunology and Histology, Semey Medical University, Semey, Kazakhstan

⁴Department of Biotechnology and Microbiology, Faculty of Natural Sciences, L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan

(*Corresponding author's e-mail: oiz5@yandex.ru)

Received: 6 November 2020, Revised: 19 July 2021, Accepted: 26 July 2021

Abstract

We studied the combined effect of gamma radiation (6 Gy, remote period) and immobilization stress on immunological reactivity. Materials and methods. To address this, we carried out experiments on 40 white male Wistar rats weighing 220 ± 20 g, which were divided into 4 groups: Group I - intact animals, group II - animals exposed to gamma radiation (dose of 6 Gy), group III - animals exposed to immobilization stress and group IV - animals exposed to combined exposure to gamma radiation (dose of 6 Gy) and immobilization stress. Animals in groups II and IV were irradiated 90 days before the study on the TERAGAM radiotherapeutic cobalt unit. In groups III and IV, animals were modeled for acute immobilization stress by immobilization for 6 h in bright light. Results. At the early period of immobilization stress, all indicators of the immune system, such as cellular, humoral and nonspecific phagocytic links of immunity were activated, indicating the activation of the general adaptive syndrome of the organism. The suppressive action of gamma radiation on the immune system was preserved in a remote period, with this suppression being revealed not only on the T-cellular link, but also on nonspecific phagocytic links, and the functional metabolic activity of neutrophils. In the remote period after the combined effect of sublethal gamma radiation and immobilization stress in the early stage of the adaptation syndrome, there was a decrease in all cells of the T system of immunity, in the functional ability of leukocytes, and in the mononuclear phagocytic system of the body. The experimental immobilization-radiation pathological process was accompanied by disorders of the functional activity of the essential adaptive systems of the body. Based on our results, we could conclude about the dominant role of ionizing radiation in immunological reactivity.

Keywords: Radiation, Immobilization stress, Immune system, Combined action, Remote period

Introduction

The occurrence of leukemia and malignant tumors are the most often remote effects of radiation. It has also been shown that the health effects of radiation might depend on the duration of the exposure [1,2]. The damage by ionizing radiation is known to be mainly due to lipid peroxidation caused by active oxygen forms. The higher the level of the active oxygen form, the greater the oxidative damage [3,4]. In the long term, even a small dose of radiation could cause genetic changes, oncogenesis and physiological changes, associated with a compromised immune system and increased cellular stress [5,6]. In contrast, it is also believed that a low dose of radiation could induce both positive and negative bioeffects, thereby reducing carcinogenesis, increasing longevity and increasing fertility [7-9].

Absorption of energy from ionizing radiation by genetic material in the cell is known to lead to DNA damage, which in turn leads to chromosome aberrations and gene mutations. Both early and late

effects of radiation have been reported to lead to organ and tissue damage caused by cell death, which in turn leads to the development of cancer. Epidemiological studies have demonstrated a dose-response relationship for the induction of cancer [10]. For instance, aberrations of the chromosomal complex of peripheral blood lymphocytes were revealed in the long term following the incidence of leaked ionizing radiation at the Chernobyl nuclear power plant (ChNPP) [11].

The results of this study revealed a high prevalence of chronic diseases among the population living near radioactive waste storage facilities. The population was reported to also receive chronic irradiation as a consequence of the long-term exposure to radiation, with the results of the study pointing to an almost complete absence of healthy people living in the area. Diseases of the cardiovascular, hematopoietic, respiratory and musculoskeletal systems occupied the top places in the prevalence of diseases among the population exposed to radiation [12-14]. According to the results obtained from field and laboratory analytical studies, the authors identified the negative impact of the tailings dump (radioactive waste storage) of the Stepnogorsk mining and chemical plant (uranium mining enterprise, Republic of Kazakhstan) on the environment of adjacent areas, expressed as radionuclides contaminating the soil, water and vegetation. The population living in local radioactive areas was estimated to receive an annual effective dose of about 6.5 mSv/y, whereas the normal level from natural sources should be 1 mSv/y [15].

Recently, much has been said about the “untargeted effects” of ionizing radiation in a remote period. Briefly, “targetless effect” is considered the ability to transfer changes from irradiated cells (target cells for radiation exposure) to unirradiated cells. This could be considered as one of the directions of the distant effects of ionizing radiation. The authors also received new data related to the radiation “witness effect” and adaptive response. It was found that there are 2 mechanisms - intercellular interactions and secret factors that could be transmitted at a distance. The importance of studying “nontargeted effects” is related to the use of research results in theoretical radiobiology, as well as in radiation ecology, when studying the long-term consequences of human-made disasters [16,17]. It has been shown that “in radiation disasters, mankind for the first time has encountered the complex, multifactorial stress,” which includes not only the biological effects of radiation but also the psychological stress of a complex structure. The psychological manifestations and consequences of this stress have been associated with the expectation of a health catastrophe [18].

One of the most radiosensitive functions of human and animal organisms is immunological reactivity. A characteristic feature of exposure to radiation is known to be the long preservation of damage in separate parts of the immune system and its associated remote consequences and complications manifested by the development of malignant tumors [19]. Ionizing irradiation in different doses in remote periods might lead to the suppression of the innate and acquired humoral and cellular immunity. Whole-body radiation exposure at doses > 2 Gy (Gray) could cause different clinical symptoms; higher doses could be so acute that they would become life-threatening [20].

Studies on the health effects of remote exposure to ionizing radiation are relevant worldwide, including Kazakhstan; some areas have been exposed to local fallout from nuclear tests at the Semipalatinsk nuclear test site; the populations living in these areas have been exposed to internal and external exposure. People who received a specific dose of radiation while living in these areas, might also face many other daily difficulties, causing further emotional distress. Given the importance of the immune system in the formation of pathological processes, its high sensitivity, as well as the significant consequences following its compromise, we were interested in its role in the formation of pathological processes in combination with the action of stress and remote period of exposure to ionizing radiation.

Materials and methods

Subjects

To address this, we carried out experiments on 40 white male Wistar rats weighing 220 ± 20 g (Kennel laboratory animals “Pushchino”, Moscow region, Russia), which were divided into 4 groups: Group I - intact animals, group II - animals exposed to gamma radiation (dose of 6 Gy), group III - animals exposed to immobilization stress and group IV - animals exposed to combined exposure to gamma radiation (dose of 6 Gy) and immobilization stress.

Irradiation

Animals in groups II and IV were irradiated 90 days before the study on the TERAGAM radiotherapeutic cobalt unit (“ISOTREND spol. S.r.o.”, Czech Republic); the radionuclide source used was cobalt-60 (Co60). The experimental animals received a single sublethal dose of 6 Gy. All experiments on animals were conducted according to the Geneva Convention (1990), the Helsinki

declaration on humane treatment of animals, and the ethical standards of the local ethics committee of the JSC “Astana Medical University,” Kazakhstan under the protocol No. 4 of 05.09.2019.

Before irradiation, we carried out a topometric and dosimetric preparation of all experimental animals for radiation. Briefly, the object was placed on the isocentric therapeutic table of the “Terasix” (X-ray simulator, of Czech Republic), which, based on its design and parameters, corresponded to the therapeutic table of a gamma irradiator. After being displayed on screen, the pattern of irradiated animals was directly entered into the planning system via the network connection of the computer using the digitizer. We calculated the isodose with the help of the “PlanW-2000” (UJP Praha) planning system with the topometric-dosimetric map containing technical parameters and planned doses of irradiation. The animals were exposed once to a total gamma radiation dosage of 6 Gy: SSD - 97.2 cm, SAD - 100.0 cm, field 40×40 cm², t = 356 s (SSD - distance from the source of ionizing radiation in the device to the conventional center of the irradiated pathological focus; SAD - distance from the source of ionizing radiation in the device to the nearest surface of the irradiated object) [21]. During irradiation, animals were in a specially designed organic glass cell, with multiple isolated cells being employed for each animal [22].

Stress modeling

In groups III and IV, animals were modeled for acute immobilization stress by immobilization for 6 h in bright light. The animals were subjected to stress for 1 day, 3 times for 6 h with 1 h breaks (the total experiment lasted for 20 h). For this, experimental animals were placed in individual plastic compartments adapted for immobilization (length - 20 cm, width - 6 cm). The animals were euthanized 1 h after the induction of acute immobilization stress by incomplete decapitation under mild ether anesthesia.

Immunofluorescence, phytohemagglutinin and immunodiffusion

Blood was taken in heparin tubes (25 units/mL) to assess their immunological status. Accordingly, we determined the total number of leukocytes and lymphocytes in the peripheral blood of all experimental animals. The number of CD19+ and CD3+ lymphocytes and their subpopulations were determined by immunofluorescent cell staining using antibodies conjugated with FITC, explicitly adapted for experiments with rats. CD3+, CD4+, CD8+ and CD19+ FITC monoclonal antibodies were acquired from GALTAG Laboratories (5791 Van Allen Way, Carlsbad, CA 92008 USA). The stained cells were examined with a fluorescent microscope (EVOSTM FL, Thermo Fisher Scientific). Surveillance was conducted in a darkened room. We determined both the inhibition reaction of leukocyte migration (LMIR) with phytohemagglutinin (PHA-M, GibcoTM) [23], as well as the concentration of circulating immune complexes (CIC) [24]. We determined the content of immunoglobulins A, M, G, using radial immunodiffusion on Mancini agar gel [25]. We also carried out the determination of the phagocytic activity (PA) of neutrophils [26] and performed nitro blue tetrazolium tests (NBT) [27].

Statistics

All the results were presented as mean ± standard error of mean (SEM) using GraphPad Prism (GraphPad Software version 5.01, Inc., USA). Statistical analyses with *p*-value less than 0.05 were considered significant.

Results and discussion

The immune system in irradiated animals at a dose of 6 Gy

Our study showed that exposure of experimental animals to a high dose of radiation in a remote period acted as an immunosuppressive agent, revealing a significant sensitivity of leukocytes and T-lymphocytes and their subpopulations. Hence, we observed the occurrence of leukopenia in exposed animals; the number of leukocytes was reduced by 18.72 % (*p* < 0.05) (**Table 1**).

At the same time, both the percentage and the absolute number of lymphocytes were shown to be increased by 8.58 and 5.74 % (*p* < 0.05), respectively. Regarding T lymphocytes, we identified the following changes: the total number of lymphocytes was reduced by almost 45.03 % (*p* < 0.001), and the relative number by 35.12 % (*p* < 0.05), leading to a significant reduction of cells with CD4+ helper activity: The total number was reduced by 26.32 % (*p* < 0.05), and the relative number by 30.27 % (*p* < 0.05). Thus, the immunoregulatory index (IRI = CD4+/CD8+) in animals exposed to radiation was reduced in comparison with the intact group. In the case of T-suppressors, an inhibition of their activity was observed, also exhibiting a decrease in the total and relative number by 22.22 (*p* < 0.05) and 36.86 %

($p < 0.01$), respectively. It is known that the activity of the CD3+ lymphocyte lymphokine-producing cells reflects the functional activity of the T system of immunity. Our study revealed a decrease in the production capacity of CD3+ lymphocytes in irradiated animals, as well as a 32.56 % increase in the LMIR migration index in the PHA-treated compared with the control group ($p < 0.05$). We found that the ability of cells to produce cytokines suppressing the migration of leukocytes was expressed in the control group. Whereas, in animals under the influence of gamma radiation, the production of cytokines was reduced, thus indicating the immunosuppressive action of radiation, which led to the suppression of the synthesis of cytokines, as the migration index was higher than that in intact animals. According to several studies [28,29], ionizing radiation could cause a decrease in DNA synthesis and the number of cells transformed into blasts under the influence of PHA, reducing the production of the factor inhibiting the migration of macrophages (MIF) and the associated cytotoxic effect.

Against the background of the explicit suppression of cell immunity, we obtained the result of the immunoregulatory index (CD4+/CD8+, IRI), exhibiting a shift to the left towards a decrease. The decrease in IRI is considered typical in immunodeficiency states and cancers. Thus, the pathognomonic laboratory symptom of immune deficiency was shown to gradually include incomplete numbers of CD4+ and a decrease of IRI. The decrease observed in IRI in our example was assumed to be due to a sharp decrease in the number of T cells with helper activity compared with those of T cells with suppressor activity. As is known, CD4+ cells stimulate the CD19+ lymphocyte population for production of antibodies. Insufficient helper function of T-lymphocytes might lead to a decrease in the sensitivity of the body to antigenic stimulation, contributing to the development of severe infectious complications and pathological conditions of radiation genesis.

Taking into account the complexity of the character of changes in the humoral immunity with the radiation lesion of the organism, we aimed to study the long-term reaction of humoral immunity in interrelation with the nonspecific phagocytic link of immunity at the same dose of 6 Gy gamma-irradiation. To determine the state of humoral and nonspecific phagocytic links of immunity in the case of gamma irradiation, we studied the number of CD19+ lymphocytes, the concentration of CIC, serum immunoglobulins, phagocytic activity, phagocytic number and also performed a NBT-test.

The suppressed levels of CD4+ cells, as well as the persistent level of CD19+ cells indicated the implementation of the immune response, which, thus, activated the B-cells; proof of this was a reliable increase in the absolute and relative number of CD19+ lymphocytes in the blood by 37.5 and 41.99 %, respectively compared with control ($p < 0.05$). B lymphocytes, in turn, are known to switch from IgM synthesis to IgG and IgA synthesis as they mature, but in this case we observed a suppression in the levels of IgG and IgA, whereas the concentration of IgM was shown to be increased by 43.11 % ($p < 0.05$) in irradiated animals. The decrease of IgA (by 16.29 %, $p < 0.05$) and IgG (by 32.66 %, $p < 0.01$) was an important criterion of the immunosuppressive action of radiation in humoral immunity. B-lymphocytes carry fixed monomeric immunoglobulin molecules on their outer membranes, which are used as antigen-recognition receptors. Plasma cells are capable of producing immunoglobulins of only 1 class (IgA, IgM, IgG, IgD and IgE). The choice of the class of immunoglobulins synthesized by the plasma cell is determined by T-helpers. There is a slowdown in the process of switching B-lymphocytes from IgM to other immunoglobulins synthesis. Perhaps, this is due to a deficiency of T-lymphocytes (Th1), which, if in sufficient numbers, induce the switching of B-lymphocytes from the synthesis of immunoglobulin of one class to another. The main factor in the suppression of T-helpers in our case is ionizing radiation. Irradiation had also a significant impact on the concentrations of CIC. As such, we identified a reliable decrease of 21.64 % ($p < 0.05$) in the blood serum of experimental rats.

Serious changes were observed in the nonspecific phagocytic immunity link in irradiated organisms. We observed a reduction of the phagocytic activity of blood cells by 17.31 % ($p < 0.05$) in irradiated experimental animals. The results of the calculation of the average number of latex absorbed by 1 phagocyte (phagocytic number) in the group of irradiated animals revealed a decrease by 9.35 % ($p < 0.05$).

Evaluating the oxygen-dependent phagocytic killing in the recovery test of NBT is known to serve as an indicator of the phagocytic and metabolic activity of neutrophilic granulocytes. A NBT-test might reflect the final reaction of one of the critical enzymatic systems responsible for the effector potential of phagocytes. Our study showed that the index of the NBT-test in experimentally irradiated animals was decreased by 13.12 % ($p > 0.05$), indicating a suppression of the functional activity of neutrophils. Collectively, the revealed tension of the phagocytic link in the form of the suppression of the phagocytic activity, reduction of the phagocytic number and obtained NBT-test results in irradiated animals further testified for the immunodepressant effect of radiation.

Thus, we revealed a number of changes in experimental rats irradiated with gamma radiation, which were characterized, first of all, by a decrease of the absolute and percentile number of T lymphocytes and their subpopulations (CD4+ and CD8+), a decrease in the functional activity of T-lymphocytes, and a decrease in the protective and adaptive mechanisms of the organism in response to the action of long term exposure to a high dose of radiation. The decrease in the number of CD3+ cells and their subpopulations, which are known to serve as the response of the immune system to developing pathological processes, could be regarded as a general physiological response of an organism in response to a stress irritant [30]. However, there were also changes in qualitative and quantitative indicators of humoral and nonspecific phagocytic links of immunity, which constitute a fact of developed immunodeficiency of radiation genesis. Accordingly, we found that the proliferative activity of T-cells, and quantitative and functional states of the T system of immunity were also decreased.

The analysis of the factual material showed that the suppressive action of gamma radiation on the immune system was preserved in the long term; suppression was detected not only on the T-cell link, but also on nonspecific phagocytic links and on the functional metabolic activity of neutrophils.

The immune system under immobilization stress

We found that the number of leukocytes in animals under immobilization stress (withdrawn from the experiment after 1 h of immobilization stress) was increased by 41.49 % ($p < 0.05$) compared with the intact group (**Table 1**). Furthermore, the total number of lymphocytes in the animals of group III was increased by 60.07 % ($p < 0.01$), as did the relative number of lymphocytes ($p > 0.05$). The absolute number of CD3+ lymphocytes in experimental animals was increased by 1.18 times ($p < 0.05$), whereas the percentage of lymphocytes remained within the levels of those in the intact group ($p > 0.05$). Regarding the CD4+ cells, their total number was shown to be increased by 31.57 % ($p < 0.05$) in group III after 1 h of stress, with their relative number also exhibiting an increasing tendency ($p > 0.05$). In the case of CD8+ lymphocytes, a similar picture was revealed: Increase of their absolute number by 24.07 % ($p < 0.05$) and relative number by 26.90 % ($p < 0.05$). We also noted an increase of IRI in group III by 6.42 % ($p > 0.05$). We also identified an increase of the production capacity of CD3+ lymphocytes and a 24.42 % decrease of the LMIR migration index in PHA in experimental animals in comparison with the intact group ($p < 0.05$). Thus, it was suggested that the production of cytokines was stimulated on the background of stress.

To determine the state of humoral and nonspecific phagocytic links of immunity under stress, we evaluated the number of CD19+ lymphocytes in peripheral blood, the concentration of CIC, immunoglobulin in serum, and determined phagocytosis, phagocytic number, as well as performed a NTT-test. Our results showed that 1 h after stress, the absolute and relative number of CD19+ lymphocytes in the blood of animals increased by 1.65 and 1.91 times, respectively, compared with that in the control ($p < 0.001$). The serum concentration of CIC was observed to be decreased from 1.34 ± 0.04 to 0.71 ± 0.07 g/L ($p < 0.001$). Against the background of the activation of cellular immunity in the investigated animals, we observed the increase in the levels of IgA and IgG: The concentration of IgA increased by 60 % ($p < 0.01$), while that of IgG by 88.52 % ($p < 0.001$), demonstrating the importance of these indicators for assessing the level of humoral immunity under stress.

In animals exposed to immobilization stress, the phagocytic activity of blood cells was shown to be increased by 64.32 % ($p < 0.01$), with the phagocytic number being increased by 82.01 % ($p < 0.01$). We also found that the NTT-test index was increased by 76.67 % ($p < 0.01$), indicating the activation of the functional activity of neutrophils.

Thus, we demonstrated here the changes that occurred in humoral immunity of experimental rats undergoing immobilization stress, which were primarily characterized by an increase in the number of white blood cells, the absolute and percentile of CD3+ lymphocytes and their subpopulations (CD4+ nad CD8+), as well as an increase in the functional activity of CD3+ lymphocytes and the protective mechanisms of the body in response to stress. The currently conducted study exhibited the occurrence of activation in all indexes, both in cellular, humoral, and nonspecific phagocytic links of immunity, and thus activation of the general adaptive syndrome of an organism at the early period of immobilization stress.

The immune system under combined exposure to gamma radiation (dose of 6 Gy, remote period) and immobilization stress

Our study showed that combined exposure of experimental animals to a high dose of radiation in the remote period and immobilization stress acted as an immunosuppressive agent, revealing the dominant

role of ionizing radiation. We noted the occurrence of leukopenia in experimental animals, with their number of leukocytes being reduced by 19.65 % ($p < 0.05$) (Table 1).

Table 1 Immune system indices in the blood of experimental animals under the influence of stressogenic factors, $M \pm m$.

Exponents	Group I - intact animals	Group II - animals exposed to gamma radiation (dose of 6 Gy)	Group III - animals exposed to immobilization stress	Group IV - animals exposed to combined exposure to gamma radiation and immobilization stress
Leucocytes, $\times 10^9/L$	6.41 ± 0.17	$5.21 \pm 0.34^*$	$9.07 \pm 0.75^*$	$5.15 \pm 0.33^*$
Lymphocyte, $\times 10^9/L$	2.68 ± 0.11	2.91 ± 0.21	$4.29 \pm 0.32^{**}$	$2.24 \pm 0.17^*$
Lymphocyte, %	37.81 ± 3.31	35.64 ± 2.45	44.37 ± 3.24	40.22 ± 3.75
CD3+	Abs.n. ^o	1.51 ± 0.08	$0.83 \pm 0.05^{***}$	$1.79 \pm 0.11^*$
	%	30.61 ± 2.53	$19.86 \pm 1.24^*$	33.15 ± 2.37
CD4+	Abs.n. ^o	0.76 ± 0.06	$0.56 \pm 0.04^*$	$1.00 \pm 0.08^*$
	%	19.72 ± 1.57	$13.75 \pm 0.71^*$	21.45 ± 1.65
CD8+	Abs.n. ^o	0.54 ± 0.04	$0.42 \pm 0.03^*$	$0.67 \pm 0.05^*$
	%	11.04 ± 0.76	$6.97 \pm 0.53^{**}$	$14.01 \pm 1.25^*$
CD19+	Abs.n. ^o	0.40 ± 0.03	$0.55 \pm 0.03^*$	$0.66 \pm 0.04^{***}$
	%	6.81 ± 0.54	$9.67 \pm 0.73^*$	$13.07 \pm 1.07^{***}$
CD4+/CD8+	1.40 ± 0.10	1.33 ± 0.08	1.49 ± 0.12	1.30 ± 0.11
CIC, y.e.	1.34 ± 0.04	$1.05 \pm 0.08^*$	$0.71 \pm 0.07^{***}$	$0.99 \pm 0.08^*$
ITML	0.86 ± 0.05	$1.14 \pm 0.08^*$	$0.65 \pm 0.05^*$	$1.11 \pm 0.09^*$
Phagocytes, %	36.10 ± 2.46	$29.85 \pm 1.87^*$	$59.32 \pm 4.36^{**}$	$27.52 \pm 2.51^*$
Cytophagous number	1.39 ± 0.11	1.26 ± 0.08	$2.53 \pm 0.20^{**}$	1.21 ± 0.13
NTT-test, %	4.80 ± 0.37	4.17 ± 0.22	$8.48 \pm 0.64^{**}$	4.08 ± 0.31
Ig A g/L	5.71 ± 0.40	$4.78 \pm 0.21^*$	$9.14 \pm 0.72^{**}$	4.86 ± 0.32
Ig M g/L	6.82 ± 0.43	$9.76 \pm 0.73^*$	6.26 ± 0.47	$9.56 \pm 0.67^*$
Ig G g/L	7.93 ± 0.53	$5.34 \pm 0.37^{**}$	$14.95 \pm 1.02^{***}$	$5.68 \pm 0.42^*$

Note: Changes statistically significant in comparison with intact control * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. ^o absolute number of cells $\times 10^9/L$.

A significant reduction was shown in the absolute number of lymphocytes in blood from $2.68 \pm 0.11 \times 10^9/L$ to $2.24 \pm 0.17 \times 10^9/L$ ($p < 0.05$), whereas their relative number was not markedly changed ($p > 0.05$).

The related content of CD3+ lymphocytes in experimental rats with an absolute reduction in their number by 48.34 % ($p < 0.001$), was demonstrated to correspond to that of the intact value. The subpopulation of T lymphocytes under these conditions exhibited the following changes: The absolute number of CD4+ was decreased by 38.16 % ($p < 0.01$), the relative by 28.9 % ($p < 0.05$), the absolute number of CD8+ by 33.33 % ($p < 0.05$) and the relative by 33.24 % ($p < 0.05$). At the same time, the immunoregulatory index was shown to be accordingly decreased due to a reduction of the number of lymphocytes with helper activity. During this period, we noted a reliable increase in the index of leukocyte migration in the LMIR reaction to PHA from 0.86 ± 0.05 to 1.11 ± 0.09 , indicating a decrease in the functional activity of the T-system of immunity ($p < 0.05$).

Collectively, our results indicated that following the combined effect of remote period sublethal gamma radiation and immobilization stress, there was a reduction of the total number of lymphocytes; this decrease in some CD3+ and several CD4+ lymphocytes, which appeared to play a considerable role in immunological reactivity, was also reflected in the low indicators of IRI. In addition, we found that in parallel with a decrease in the number of CD8 + lymphocytes, the functional capacity of T-lymphocytes was also decreased.

Our study results showed that under the combined effect, the absolute and relative number of CD19+ lymphocytes in the blood of animals was increased by 20 and 48.46 %, respectively, compared with the control ($p < 0.05$). This combination had also a significant impact on the concentrations of CIC. Thus, we observed a 26.12 % decrease ($p < 0.05$) in the concentration of CIC in the serum of experimental rats. At this background of the increase in the concentration of CD19+ in investigated animals, we identified a suppression in the levels of IgA (14.89 %, $p > 0.05$) and IgG (23.37 %, $p < 0.05$), whereas the concentration of IgM was shown to be increased to 40.17 % ($p < 0.05$). The same result was obtained during isolated radiation exposure, thereby detecting the radiation trace of the effect on the humoral immunity that explains the pathogenetic causes of the increase in IgM and suppression of IgA and IgG levels. Regarding the phagocytic activity of blood cells, we noted a suppression by 23.77 % ($p < 0.05$) and a decrease in the phagocytic number by 12.95 % ($p > 0.01$).

Moreover, the NBT-test indicator exhibited a downward trend. It should be noted that the implementation of recovery processes in the body is known to be facilitated in the remote period after exposure with a decrease in the dose rate. Under irradiation of the organism with different power, the degree of the response of its reserve adaptive capacity would be different. The revealed changes in the remote period characterizing the immunodepression of the irradiated organism could be considered as an adaptive compensatory reaction of the organism in response to radiation alone and in combination with nonradiative factors [31].

Thus, in the remote period after the combined exposure to sublethal gamma irradiation and immobilization stress in the early stage of the adaptation syndrome, all cells of the T system of immunity, such as CD4+/CD8+, as well as the functional ability of leukocytes and mononuclear phagocytic system of the organism decrease. Moreover, we noted a significant depression of immunoglobulins A and G, whereas IgM was shown to be activated. The decrease in the content of IgA and G testified to the occurrence of immunodeficiency, which is known to be accompanied by chronic inflammation and the presence of high tendency for the development of neoplasm at high doses of ionizing radiation. Based on the results of this study, we could conclude about the dominant role of ionizing radiation in the immunological reactivity of organisms.

Conclusions

The effect of remote period gamma radiation remained suppressive on the immune system, including nonspecific phagocytic links, and the functional metabolic activity of neutrophils. Hence, the mechanisms of development of stress response to radiation genesis might depend on the adaptation and insufficiency of physiological measures for the protection of the irradiated organism. At the early period of immobilization stress, all indexes of the immune system, such as cellular, humoral and nonspecific phagocytic links of immunity were activated, indicating the activation of the general adaptive syndrome of the organism. In the remote period after the combined action in the early stage of the adaptation syndrome, there was a decrease in all cells of the T system of immunity, in the functional ability of leukocytes and in the mononuclear phagocytic system of the organism. Ionizing radiation, combined with immobilization stress had a more pronounced effect on the formation of immunodeficiency syndrome in a remote period. Moreover, the dominant agent affecting the developing pathology under the combined effects of the above-mentioned 2 factors was shown to be ionizing radiation.

Acknowledgments

The work was supported by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan, grant No. 0118RK00662.

References

- [1] TV Azizova, NG Semenikhina and MB Druzhinina. Multi-organ involvement and failure in selected accident cases with acute radiation syndrome observed at the Mayak Nuclear Facility. *Br. J. Radiol.* 2005; **78**, 30-5.
- [2] T Verbiest, R Finnon, N Brown, L Cruz-Garcia, P Finnon, G O'Brien, E Ross, S Bouffler, CL Scudamore and C Badie. Tracking preleukemic cells *in vivo* to reveal the sequence of molecular events in radiation leukemogenesis. *Leukemia* 2018; **32**, 1435-44.
- [3] Y Kojima, T Kondo, QL Zhao, M Shoji and R Futasuya. Protective effects of cimetidine on radiation-induced micronuclei and apoptosis in human peripheral blood lymphocytes. *Free Radic. Res.* 2002; **36**, 255-63.
- [4] EI Azzam, JP Jay-Gerin and D Pain. Ionizing radiation-induced metabolic oxidative stress and prolonged cell injury. *Canc. Lett.* 2012; **327**, 48-60.
- [5] N Shimura and S Kojima. Effects of low-dose-gamma rays on the immune system of different animal models of disease. *Dose Response* 2014; **12**, 429-65.
- [6] A Graupner, DM Eide, C Instanes, J Andersen, D Brede, S Dertinger, O Lind, A Brandt-Kjelsen, H Bjerke, B Salbu, D Oughton, G Brunborg and A Olsen. Gamma radiation at a human relevant low dose rate is genotoxic in mice. *Sci. Rep.* 2016; **6**, 32977.
- [7] FR Tang, WK Loke and BC Khoo. Low-dose or low-dose-rate ionizing radiation - induced bioeffects in animal models. *J. Radiat. Res.* 2017; **58**, 165-82.
- [8] RK Kim, MJ Kim, KM Seong, N Kaushik, Y Suh, KC Yoo, YH Cui, Y Jin, S Nam and SJ Lee. Beneficial effects of low dose radiation in response to the oncogenic KRAS induced cellular transformation. *Sci. Rep.* 2015; **5**, 15809.
- [9] JM Cuttler. Health effects of low level radiation. When will we acknowledge the reality? *Dose Response* 2007; **5**, 292-8.
- [10] K Suzuki and S Yamashita. Low-dose radiation exposure and carcinogenesis. *Jpn. J. Clin. Oncol.* 2012; **42**, 563-8.
- [11] AM Nikiforov, NV Makarova and NN Zybyna. Chromosomal aberrations and indicators of oxidative stress in liquidators of the consequences of the Chernobyl accident in the remote period after the action of ionizing radiation. *Cytology* 2004; **6**, 561-6.
- [12] Y Shimizu, K Kodama, N Nishi, F Kasagi, A Suyama, M Soda, E Grant, H Sugiyama, R Sakata, H Moriwaki, M Hayashi, M Konda and R Shore. Radiation exposure and circulatory disease risk: Hiroshima and Nagasaki atomic bomb survivor data, 1950 - 2003. *BMJ* 2010; **340**, b5349.
- [13] SC Darby, M Ewertz, P McGale, AM Bennet, UB Goldman, D Bronnum, C Correa, D Cutter, G Gagliardi, B Gigante, MB Jensen, A Nisbet, R Peto, K Rahimi, C Taylor and P Hall. Risk of ischemic heart disease in women after radiotherapy for breast cancer. *New Engl. J. Med.* 2013; **368**, 987-98.
- [14] DJ Janavayev, YT Kashkinbayev, KB Ilbekova, YA Saifulina, MM Bakhtin, MK Sharipov and PK Kazymbet. Health status of the population living in the zone of influence of radioactive waste repositories. *Electron. J. Gen. Med.* 2019; **16**, em176.
- [15] P Kazymbet, M Bahtin, E Kashkinbaev, D Dzhanabaev, Z Dzhaubayeva and M Sharipov. Radiation situation at the tailing dump of the Stepnogorsk mining-chemical combine and adjacent territories. Message I. *Meditinskaya Radiologiya I Radiatsionnaya Bezopasnost* 2018; **63**, 40-7.
- [16] II Pelevina, VV Petushkova, VA Biryukov, AV Akleyev, EA Neifakh, NGMinaeva, OV Ktitorova, AV Aleshchenko and RI Pleshakova. The "radiation-induced non-targeted effects" and their role in human cell response to low radiation forcing. *Radiat. Biol. Radioecol.* 2019; **3**, 261-73.
- [17] VN Ivanov and TK Hei. A role for TRAIL/TRAIL-R2 in radiation-induced apoptosis and radiation-induced bystander response of human neural stem cells. *Apoptosis* 2014; **19**, 399-413.
- [18] OI Kravtsov. Psychohygiene and psychoprophylaxis in the activities of specialists who take part in the elimination of the consequences of radiation accidents. *Hyg. Sanit.* 2017; **96**, 900-3.
- [19] H Gao, Z Dong, X Gong, J Dong, Y Zhang, W Wei, R Wang and S Jin. Effects of various radiation doses on induced T-helper cell differentiation and related cytokine secretion. *J. Radiat. Res.* 2018; **59**, 395-403.
- [20] International Atomic Energy Agency. *Radiation biology: A handbook for teachers and students.* International Atomic Energy Agency, Vienna, Austria, 2010.
- [21] MA Sandybaev, BA Zhetpisbayev, NA Bazarbayev and OZ Ilderbayev. Method of topometric-dosimetric preparation of experimental animals for irradiation, Republic of Kazakhstan. *Innovat. Patent Bull.* 2009; **2009**, 21845.

- [22] BA Zhetpisbayev, MN Sandybaev, OZ Ilderbayev and NA Bazarbayev. Cell for irradiation of experimental animals. Republic of Kazakhstan. *Innovat. Patent Bull.* 2009; **2009**, 21532.
- [23] AG Artemova. The phenomenon of the inhibition of migration of blood leukocytes in guinea pigs with delayed hypersensitivity to a foreign tissue agent. *Bull. Exp. Biol. Med.* 1973; **10**, 67-71.
- [24] JJ Grinkevich and AN Alferov. Determination of immune complexes in the blood of cancer patients. *Laboratornoe Delo* 1981; **8**, 493-5.
- [25] G Mancini, JP Vaerman, AO Carbonara and JF Heremans. *A single radial diffusion method for the immunological quantitation of proteins.* In: H Peeters (Ed.). *Protides of biological fluids*, Elsevier, Amsterdam, Netherlands, 1964, p. 370-3.
- [26] EA Kost and IV Stenko. *Investigation of phagocytosis by clinical and laboratory research methods.* Moscow, Russia, 1968, p. 78-80.
- [27] BS Nagoev and MG Shubich. The Value of the nitrosine tetrazolium recovery test for studying the functional activity of leukocytes. *Laboratornoe Delo* 1981; **4**, 195-8.
- [28] K Khan, S Tewari, NP Awasthi, SP Mishra, GR Agarwal, M Rastogi and N Husain. Flow cytometric detection of gamma-H2AX to evaluate DNA damage by low dose diagnostic irradiation. *Med. Hypotheses* 2018; **115**, 22-8.
- [29] J Rak, L Chomicz, J Wicz, K Westphal, M Zdrawowicz, P Wityk, M Zyndul, S Makurat and L Golon. Mechanisms of damage to DNA labeled with electrophilic nucleobases induced by ionizing or UV radiation. *J. Phys. Chem. B* 2015; **119**, 8227-38.
- [30] Y Shen, X Jiang, L Meng, Ch Xia, L Zhang and Y Xin. Transplantation of bone marrow mesenchymal stem cells prevents radiation-induced artery injury by suppressing oxidative stress and inflammation. *Oxid. Med. Cell Longev.* 2018; **2018**, 5942916.
- [31] BA Zhetpisbayev, AA Alimbayeva and KS Adrisova. Long-term effects after fractionated gamma radiation and immobilization stress on non-specific phagocytic resistance of the body. *Sci. Health* 2014; **4**, 59-62.